

Balancing the kinetochore ledger

Stuart Cane^{1,2} and Thomas J. Maresca^{1,2}

¹Biology Department and ²Molecular and Cellular Biology Program, University of Massachusetts Amherst, Amherst, MA 01003

Reduction of polo-like kinase-1 (Plk1) at kinetochores as cells progress from prometaphase to metaphase is surprising given that the kinase is thought to stabilize kinetochore–microtubule (kt–MT) attachments. In this issue, Liu et al. (2012. *J. Cell Biol.* doi:10.1083/jcb.201205090) demonstrate that kinetochore-associated Plk1 is a potent suppressor of microtubule plus-end dynamics. The authors propose that Plk1 activity facilitates the establishment of kt–MT attachments in prometaphase by stabilizing microtubules and that reduction of the kinase in metaphase promotes force generation by dynamic microtubules.

What do the kinetochore and Goldman Sachs have in common? This may sound like the beginning of a bad joke told over cocktails at an ASCB meeting, but the truth is, they have more in common than you might think. In fact, emerging research has revealed that the kinetochore carries out a balancing act rivaling that of a Wall Street banker dealing with a ledger full of complex financial derivatives.

The kinetochore ensures that the genome is accurately segregated during cell division by accomplishing two major tasks. First, it mediates the attachment of chromosomes to dynamic spindle microtubules. Second, when erroneous kt–MT attachments arise, the kinetochore fulfills its spindle assembly checkpoint (SAC) responsibilities by catalyzing production of a wait-anaphase signal that delays cell division to provide time to correct the errors. As it performs these functions, the kinetochore must manage a shifting balance of microtubule stabilizing and destabilizing activities as well as checkpoint regulators to establish and maintain kt–MT attachments and to coordinate SAC signaling.

Establishing and maintaining correct kt–MT attachments requires a precise balance between two antagonistic inputs: one favoring microtubule plus-end dynamics and the other favoring plus-end stability (reviewed by Bakhoum and Compton, 2012). Stabilizing inputs gain the upper hand over those favoring microtubule plus-end dynamics as chromosomes align during the progression from prometaphase to metaphase. Mounting evidence suggests that the transition in kt–MT attachment stability is mediated by changes in the molecular composition and phosphorylation profile of the kinetochore.


Plk1 has been implicated in both stabilizing and destabilizing kt–MT attachments. In support of a stabilizing role for

Plk1, it has been shown that disrupting Plk1 activity by either RNAi or chemical inhibition prevents the formation of stable kt–MT attachments (Sumara et al., 2004; Hanisch et al., 2006; Peters et al., 2006; Lénárt et al., 2007) and that phosphorylation of the checkpoint kinase BubR1 by Plk1 is required for normal kt–MT attachment stability (Elowe et al., 2007). On the other hand, Plk1 activity is required for the recruitment and activation of two known destabilizing factors at prometaphase kinetochores: the microtubule depolymerizing motor Kif2b (Hood et al., 2012) and Aurora B kinase (Chu et al., 2011; Moutinho-Santos et al., 2012). Depletion of Plk1 in *Drosophila melanogaster* cells leads to hyper-stable syntelic attachments (Moutinho-Santos et al., 2012)—erroneous attachments in which both members of a kinetochore pair attach to microtubules from the same spindle pole—and this finding also cuts in favor of an attachment-destabilizing function for Plk1. Moreover, inhibiting Plk1 eliminates the attachment instability otherwise generated by depletion of the phosphatase B56-PP2A (Foley et al., 2011), suggesting, again, that Plk1 activity contributes to kt–MT attachment destabilization. In this issue, Liu et al. (2012) advance our understanding of the attachment-stabilizing role of Plk1 by providing convincing evidence that Plk1 activity at the kinetochore contributes to the establishment of kt–MT attachments by suppressing microtubule plus-end dynamics.

Liu et al. (2012) tethered a fluorescence resonance energy transfer (FRET)-based Plk1 phosphorylation sensor to HeLa cell kinetochores and found, consistent with previous results showing a decline in kinetochore-associated Plk1 from pro-metaphase to metaphase (Lénárt et al., 2007), that phosphorylation of the FRET probe was reduced as chromosomes aligned. The reduction in Plk1 levels and in FRET probe phosphorylation is a result, in part, of recruitment of protein phosphatase 1 (PP1) to metaphase kinetochores, where PP1 likely dephosphorylates (and thereby renders unavailable) potential binding sites for Plk1's polo box domain (PBD). To better define the effects of Plk1 activity on kt–MT attachment stability, the authors fused a constitutively active form of Plk1 (T210D mutant) to the outer kinetochore protein Hec1 to maintain constitutively high Plk1 activity at kinetochores in metaphase. Cells expressing the Hec1-Plk1^{T210D} fusion exhibited a dramatic reduction in microtubule dynamics at the kt–MT interface compared with wild-type control cells. Suppression of plus-end dynamics by

Correspondence to Thomas J. Maresca: tmaresca@bio.umass.edu

Figure 1. Dynamic regulation of kt-MT attachment stability. Changes in the molecular composition, phosphorylation state, and structure of the kinetochore from prometaphase to metaphase mediate the establishment and maintenance of kt-MT attachments. High levels of kinetochore-associated Plk1 in prometaphase suppress microtubule plus-end dynamics, counteracting the attachment-destabilizing activity of Aurora B and facilitating the initial establishment of kt-MT attachments. The reduction of kinetochore-associated Plk1 in metaphase, mediated in part by PP1, relieves suppression of microtubule plus-end dynamics, thereby allowing dynamic kt-MTs to position attachment factors beyond the influence of Aurora B through introduction of intrakinetoche stretch. Like Plk1, kinetochore levels of B56-PP2A, CENP-E, BubR1, and Dynein, all of which have been implicated in stabilizing kt-MT attachments (Putkey et al., 2002; Lampson and Kapoor, 2005; Elowe et al., 2007; Varma et al., 2008; Foley et al., 2011), decrease upon establishment of kt-MT attachments (Hoffman et al., 2001; Foley et al., 2011). To the contrary, numerous stabilizing inputs, which include intrakinetoche stretch (Maresca and Salmon, 2009; Uchida et al., 2009), PP1 (Liu et al., 2010), Astrin-SKAP (Schmidt et al., 2010), and the Ska complex (Chan et al., 2012), increase in metaphase. CLASPI acts as part of a molecular switch that destabilizes microtubules in prometaphase with Kif2b and stabilizes microtubules in metaphase with Astrin (Manning et al., 2010).



Hec1-Plk1^{T210D} caused a metaphase arrest with reduction in both inter- and intrakinetoche stretch, accompanied by a higher incidence of merotelic attachments in which a single kinetochore attached to microtubules from both spindle poles. The authors reasoned that Plk1 is normally cleared from metaphase kinetoche to allow dynamic microtubules to exert pulling forces on the kinetochore that, in combination with PP1 activity, overcome the attachment-destabilizing effects of centromere-based Aurora B kinase. Finally, the authors considered whether Plk1 acts as a counterweight to kinetochore-associated destabilizing activities to facilitate the establishment of kt-MT attachments in prometaphase. Indeed, overexpressing the PBD, which displaces the endogenous kinase from kinetoche, significantly disrupted the establishment of kt-MT attachments after a nocodazole washout.

One fascinating subplot to emerge from this study concerns the SAC. The metaphase arrest in Hec1-Plk1^{T210D} cells was SAC dependent, as Hec1-Plk1^{T210D} cells had seven times as many Mad2-positive kinetoche as control cells, and the metaphase arrest could be overridden by chemically inhibiting the checkpoint kinase Mps1. Strikingly, the SAC-dependent arrest occurred in the presence of properly aligned chromosomes with highly stable kt-MT attachments. What might be causing this arrest? One possibility is that the cold stability and photoactivation assays deployed by Liu et al. (2012) to probe kt-MT attachment stability are not sensitive enough to detect unattached kinetoche in the Hec1-Plk1^{T210D} cells. However,

these assays are the current gold standard in the field for investigating kt-MT attachment stability, and it is difficult to imagine that they would fail to detect a sevenfold increase in unattached kinetoche. Alternatively, constitutive Plk1 activity may drive recruitment of checkpoint proteins to aligned and stably attached kinetoche. Yet, Mad2 was not detected at every Hec1-Plk1^{T210D} kinetochore, and Plk1 activity is not required for Mad2 localization (Sumara et al., 2004; Hanisch et al., 2006; Lénárt et al., 2007). Thus, we favor the view that in Hec1-Plk1^{T210D} cells, the wait-anaphase signal is generated in response to suppression of kt-MT plus-end dynamics and the resulting reduction in intrakinetoche stretch rather than in response to unattached kinetoche. Indeed, a direct role for suppressed kt-MT dynamics and reduced intrakinetoche stretch in generating a wait-anaphase signal independent of defects in kt-MT attachment has previously been hypothesized (Maresca and Salmon, 2009, 2010).

It is becoming increasingly evident that balancing stabilizing and destabilizing activities at the kinetochore is a complex undertaking (Fig. 1), and the list of influences of both kinds is already a long one. Centromeric Aurora B kinase represents perhaps the most widely recognized example of an attachment destabilizer. The influence of this kinase predominates early in mitosis until tension across the centromeres/kinetoche of bioriented chromosomes moves Aurora B substrates beyond the effective range of the kinase's activity, shifting the balance of inputs toward attachment stabilization (reviewed by Maresca and

Salmon, 2010). More recently, it has been demonstrated that Aurora B inhibits the localization of the stabilizing complexes Ska (Chan et al., 2012) and Astrin–SKAP (Schmidt et al., 2010) to prometaphase kinetochores and that both complexes later confer stability by localizing to metaphase kinetochores. In addition, kinetochore-associated phosphatase activity contributes to the formation of stable kt–MT attachments, with B56-PP2A apparently playing a key role in prometaphase (Foley et al., 2011) and PP1 in metaphase (Liu et al., 2010). Looking beyond kinase and phosphatase activity, it has also recently been shown that the outer kinetochore protein CLASP1 participates in an attachment-regulating molecular switch system. It engages with two different binding partners in succession, first forming a destabilizing CLASP1–Kif2b complex in prometaphase and then a stabilizing CLASP1–Astrin complex in metaphase (Manning et al., 2010). The results reported by Liu et al. (2012) support the hypothesis that, during prometaphase, Plk1 contributes an essential stabilizing activity to counteract the destabilizing activities that would otherwise dominate. But because Plk1 has also been implicated in kt–MT destabilization, it appears that Plk1 may sit on both sides of the kinetochore ledger. Given the recent finding that Plk1 localizes both to centromeres and, separately, to kinetochores (Carmena et al., 2012), it is possible that distinct cellular populations of Plk1 differentially regulate kt–MT attachment stability. Additional work is clearly needed to address whether and how Plk1 might play seemingly contradictory roles at the kinetochore.

So, like any successful Wall Street financier, the kinetochore is a true polo enthusiast. And, although the kinetochore may have more in common with Goldman Sachs than we thought, it is far more useful to focus on the key difference: the kinetochore actually pulls off its complex balancing act. We would benefit from learning more about exactly how.

Submitted: 23 July 2012

Accepted: 31 July 2012

References

- Bakhoum, S.F., and D.A. Compton. 2012. Kinetochores and disease: Keeping microtubule dynamics in check! *Curr. Opin. Cell Biol.* 24:64–70. <http://dx.doi.org/10.1016/j.ceb.2011.11.012>
- Carmena, M., X. Pinson, M. Platani, Z. Salloum, Z. Xu, A. Clark, F. MacIsaac, H. Ogawa, U. Eggert, D.M. Glover, et al. 2012. The chromosomal passenger complex activates Polo kinase at centromeres. *PLoS Biol.* 10: e1001250. <http://dx.doi.org/10.1371/journal.pbio.1001250>
- Chan, Y.W., A.A. Jeyaprakash, E.A. Nigg, and A. Santamaría. 2012. Aurora B controls kinetochore–microtubule attachments by inhibiting Ska complex–KMN network interaction. *J. Cell Biol.* 196:563–571. <http://dx.doi.org/10.1083/jcb.201109001>
- Chu, Y., P.Y. Yao, W. Wang, D. Wang, Z. Wang, L. Zhang, Y. Huang, Y. Ke, X. Ding, and X. Yao. 2011. Aurora B kinase activation requires survivin priming phosphorylation by PLK1. *J. Mol. Cell Biol.* 3:260–267. <http://dx.doi.org/10.1093/jmcb/mjq037>
- Elowe, S., S. Hümmel, A. Uldschmid, X. Li, and E.A. Nigg. 2007. Tension-sensitive Plk1 phosphorylation on BubR1 regulates the stability of kinetochore microtubule interactions. *Genes Dev.* 21:2205–2219. <http://dx.doi.org/10.1101/gad.436007>
- Foley, E.A., M. Maldonado, and T.M. Kapoor. 2011. Formation of stable attachments between kinetochores and microtubules depends on the B56-PP2A phosphatase. *Nat. Cell Biol.* 13:1265–1271. <http://dx.doi.org/10.1038/ncb2327>
- Hanisch, A., A. Wehner, E.A. Nigg, and H.H. Siljé. 2006. Different Plk1 functions show distinct dependencies on Polo-Box domain-mediated targeting. *Mol. Biol. Cell.* 17:448–459. <http://dx.doi.org/10.1091/mbc.E05-08-0801>
- Hoffman, D.B., C.G. Pearson, T.J. Yen, B.J. Howell, and E.D. Salmon. 2001. Microtubule-dependent changes in assembly of microtubule motor proteins and mitotic spindle checkpoint proteins at PtK1 kinetochores. *Mol. Biol. Cell.* 12:1995–2009.
- Hood, E.A., A.N. Kettenbach, S.A. Gerber, and D.A. Compton. 2012. Plk1 regulates the kinesin-13 protein Kif2b to promote faithful chromosome segregation. *Mol. Biol. Cell.* 23:2264–2274. <http://dx.doi.org/10.1091/mbc.E11-12-1013>
- Lampson, M.A., and T.M. Kapoor. 2005. The human mitotic checkpoint protein BubR1 regulates chromosome-spindle attachments. *Nat. Cell Biol.* 7: 93–98. <http://dx.doi.org/10.1038/ncb1208>
- Lénárt, P., M. Petronczki, M. Steegmaier, B. Di Fiore, J.J. Lipp, M. Hoffmann, W.J. Rettig, N. Kraut, and J.M. Peters. 2007. The small-molecule inhibitor BI 2536 reveals novel insights into mitotic roles of polo-like kinase 1. *Curr. Biol.* 17:304–315. <http://dx.doi.org/10.1016/j.cub.2006.12.046>
- Liu, D., M. Vleugel, C.B. Backer, T. Hori, T. Fukagawa, I.M. Cheeseman, and M.A. Lampson. 2010. Regulated targeting of protein phosphatase 1 to the outer kinetochore by KNL1 opposes Aurora B kinase. *J. Cell Biol.* 188: 809–820. <http://dx.doi.org/10.1083/jcb.201001006>
- Liu, D., O. Davydenko, and M.A. Lampson. 2012. Polo-like kinase-1 regulates kinetochore–microtubule dynamics and spindle checkpoint silencing. *J. Cell Biol.* 198:491–499.
- Manning, A.L., S.F. Bakhoum, S. Maffini, C. Correia-Melo, H. Maiato, and D.A. Compton. 2010. CLASP1, astrin and Kif2b form a molecular switch that regulates kinetochore–microtubule dynamics to promote mitotic progression and fidelity. *EMBO J.* 29:3531–3543. <http://dx.doi.org/10.1089/emboj.2010.230>
- Maresca, T.J., and E.D. Salmon. 2009. Intrakinetochore stretch is associated with changes in kinetochore phosphorylation and spindle assembly checkpoint activity. *J. Cell Biol.* 184:373–381. <http://dx.doi.org/10.1083/jcb.200808130>
- Maresca, T.J., and E.D. Salmon. 2010. Welcome to a new kind of tension: Translating kinetochore mechanics into a wait-anaphase signal. *J. Cell Sci.* 123:825–835. <http://dx.doi.org/10.1242/jcs.064790>
- Moutinho-Santos, T., C. Conde, and C.E. Sunkel. 2012. POLO ensures chromosome bi-orientation by preventing and correcting erroneous chromosome-spindle attachments. *J. Cell Sci.* 125:576–583. <http://dx.doi.org/10.1242/jcs.092445>
- Peters, U., J. Cherian, J.H. Kim, B.H. Kwok, and T.M. Kapoor. 2006. Probing cell-division phenotype space and Polo-like kinase function using small molecules. *Nat. Chem. Biol.* 2:618–626. <http://dx.doi.org/10.1038/nchembio826>
- Putkey, F.R., T. Cramer, M.K. Morphew, A.D. Silk, R.S. Johnson, J.R. McIntosh, and D.W. Cleveland. 2002. Unstable kinetochore–microtubule capture and chromosomal instability following deletion of CENP-E. *Dev. Cell.* 3:351–365. [http://dx.doi.org/10.1016/S1534-5807\(02\)00255-1](http://dx.doi.org/10.1016/S1534-5807(02)00255-1)
- Schmidt, J.C., T. Kiyomitsu, T. Hori, C.B. Backer, T. Fukagawa, and I.M. Cheeseman. 2010. Aurora B kinase controls the targeting of the Astrin–SKAP complex to bioriented kinetochores. *J. Cell Biol.* 191:269–280. <http://dx.doi.org/10.1083/jcb.201006129>
- Sumara, I., J.F. Giménez-Abián, D. Gerlich, T. Hirota, C. Kraft, C. de la Torre, J. Ellenberg, and J.M. Peters. 2004. Roles of polo-like kinase 1 in the assembly of functional mitotic spindles. *Curr. Biol.* 14:1712–1722. <http://dx.doi.org/10.1016/j.cub.2004.09.049>
- Uchida, K.S., K. Takagaki, K. Kumada, Y. Hirayama, T. Noda, and T. Hirota. 2009. Kinetochore stretching inactivates the spindle assembly checkpoint. *J. Cell Biol.* 184:383–390. <http://dx.doi.org/10.1083/jcb.200811028>
- Varma, D., P. Monzo, S.A. Stehman, and R.B. Vallee. 2008. Direct role of dynein motor in stable kinetochore–microtubule attachment, orientation, and alignment. *J. Cell Biol.* 182:1045–1054. <http://dx.doi.org/10.1083/jcb.200710106>